

Spring Oats - A Crop Dependent on Climate and Technological Conditions

Alina Laura AGAPIE¹, Busuioc VACARIU^{1,*}, Felicia CHETAN², Alina SIMON²
¹Agricultural Research and Development Station Lovrin, 200, Principala Street, 307250, Lovrin
²Agricultural Research and Development Station Turda, 27, Agriculturii Street, Turda

*Corresponding author: busuiocvacariu@gmail.com

Abstract. The study analyzed the productivity and quality of the spring oat variety Ovidiu. The variety was cultivated within ARDS Lovrin, under non-irrigated cultivation conditions. Two experimental factors were analyzed – crop fertilization (with two graduations) and soil cultivation (with four graduations). Production was determined quantitatively at crop harvest. Regarding the main productivity elements: panicle length, number of grains/panicle, grain weight/panicle, straw length, TKW, HLW – these were obtained by analyzing a number of 50 plants for each experimental variant. The protein percentage was determined in the laboratory using the Kjeldahl method. Among the four types of soil cultivation practiced for the establishment of the spring oat crop, for the western area of Romania the most efficient is the method of preparing the germinal bed with the heavy disc. By using this method, the costs of land preparation are reduced by 73%. The economic efficiency of disc tillage also derives from the higher yields per hectare obtained compared to the other three methods, the quality of the productions, and better storage and preservation of water in the soil.

Keywords: spring oats, plowing, scarifying, disc work, fertilization.

INTRODUCTION

The exploitation of agricultural land through incomplete or incorrect strategies seriously affects both the quantity and quality not only of agricultural production but especially of soil resources.

Crop development depends on the growth and development of plant roots in loose soil with good water retention capacity and nutrient absorption. In this regard, optimal soil tillage is required, while the use of plowing methods generally leads to soil compaction (Gürsoy, 2021; Mileusnić et al., 2022). Climate change, manifested through atmospheric drought, decreased precipitation, and uneven distribution throughout the agricultural year, requires urgent measures for conserving and retaining water in the soil (Chețan, 2021; Rusu et al., 2011).

The use of alternative soil tillage methods, with minimal disturbance, is an alternative to conventional tillage; however, these methods must be thoroughly studied, considering that soil tillage is dependent on the soil type (Dinuta, 2024). Through the use of minimum tillage methods, a large portion of plant residues is incorporated or maintained on the surface, functioning as a natural mulch (Chețan et al., 2021; Copcea et al., 2018; Derpsch, 2001). High yields of spring crops, which increasingly dominate crop structures, can be ensured through appropriate fertilization and agronomic practices adapted to each crop (Bărdaș et al., 2022; Ceclan et al., 2022; Chiriță et al., 2024; Șimon et al., 2022).

Oats are ranked sixth in terms of cultivation area and productivity, after wheat, corn, rice, barley, and sorghum. The presence of adapted varieties and favorable climatic conditions allows for successful cultivation in countries such as the United Kingdom, Germany, Sweden, Finland, and France (Hoffman, 1995). It is a crop well adapted to difficult climatic conditions—low temperatures and low precipitation. To achieve high yields, optimized technology is necessary, including the appropriate sowing period (Zhang et al., 2019).

Global studies show that drought and atmospheric heat are the main causes of low cereal yields (IPCC, 2014). Abiotic stress drastically reduces cereal productivity, affecting plant growth and development, biomass accumulation, and may even cause plant death (Witcombe et al., 2008; Deryng et al., 2014; Niu et al., 2014). Climate predictions are not particularly encouraging—average temperatures continue to rise, and extreme weather events are becoming more frequent (Battisti & Naylor, 2009; Diez et al., 2012).

Given the increasingly evident climate changes, it becomes necessary to study and understand crop responses to extreme conditions. Stress tolerance is the plant's ability to grow and produce yields despite the presence of abiotic stress (Mickelbart et al., 2015). Acclimatization is one of the most important processes that cereal crops must undergo (Thomashow, 2010). The duration of this process varies from a few hours—for thermal stress (Senthil-Kumar et al., 2003; Rampino et al., 2009)—to several days for frost stress (Byun et al., 2014). During the acclimatization process, the expression level of many stress-responsive genes is altered (Hua, 2009; Jia et al., 2016).

Cereal crops, having a relatively underdeveloped root system, can be cultivated under various soil tillage conditions. This project analyzed four types of soil work (plowing, heavy disc, deep scarification, shallow scarification) for the spring oat crop, aiming to identify the optimal variant for the western region of the country, one that ensures high-quality yields with minimal disturbance to soil structure.

MATERIAL AND METHODS

The experiment was established at ARDS Lovrin on a typical, weakly gleyed, medium clay loam chernozem.

The spring oat crop, variety *Ovidiu*, was studied in relation to the type of soil tillage and fertilization system, with the following grading of experimental factors:

Factor A – type of soil tillage:

- a1 – soil tillage using a heavy disc
- a2 – soil tillage through shallow scarification (20 cm)
- a3 – soil tillage through deep scarification (50 cm)
- a4 – soil tillage through plowing (18 cm)

Factor B – crop fertilization:

- b1 – NPK 15:15:15 – 150 kg of commercial product
- b2 – NPK 15:15:15 – 150 kg of commercial product + 200 kg of commercial ammonium nitrate (33.5%).

An analysis of the precipitation regime data shows that the 2023–2024 agricultural year was an atypical one overall (Figure 1), with recorded rainfall values significantly below the multiannual monthly average in most months of the year. The uneven distribution of rainfall—by decade, month, and year—had a negative impact on crop development. The total amount of rainfall during the 2023–2024 agricultural year was 417.4 mm, which represents a negative deviation of 104 mm compared to the multiannual average. During the vegetation period of the spring oat crop (March–July), the total rainfall was 131.6 mm, with a negative deviation of 124.6 mm from the multiannual average. During this period, out of 153 days, there were 123 days with no rainfall at all, 12 days with daily rainfall between 0–1 mm, 26 days with daily rainfall between 1–10 mm, and only 2 days with daily rainfall exceeding 10 mm.

This rainfall deficit, combined with excessive temperatures and atmospheric drought, intensified the phenomenon of both soil and atmospheric drought, drastically reducing yields per unit area.




Figure 1. Precipitations evolution at ARDS Lovrin, 2023–2024

The average annual temperature was 14.6°C, which was 3.6°C higher than the multiannual average over 70 years (10.9°C) (Figure 2). During the crop growing season, monthly temperatures exceeded the multiannual average. The associated atmospheric heat, along with the lack of precipitation, negatively affected the progression of phenophases and, consequently, the yield obtained. The largest deviations from the average were recorded from February to July. Thus, in February, a positive deviation of +7.0°C was recorded, in March +4.3°C, in April +3.1°C, in June +3.8°C, and in July +4.6°C.




Figure 2. Temperature evolution at ARDS Lovrin, 2023-2024

RESULTS AND DISCUSSIONS

The results obtained under the conditions of the 2023-2024 agricultural year are analyzed and presented below. The production, the main elements of productivity and the quality of the crop were analyzed, under the various conditions imposed by the experimental factors.

In Table 1 are presented the production results.

Table 1. Table of comparisons for production (interaction soil work x fertilization)

Experimental factors		Production	Difference	%	Significant
NPK	Disc harrowing	3413,5	100,0	0,00	Control
	Light scarification	3339,0	97,8	-74,50	-
	Deep scarification	3172,0	92,9	-241,00	-
	Plowing	3227,0	94,5	-186,00	-
NPK + NH ₄ NO ₃	Disc harrowing	3834,0	100,0	0,00	Control
	Light scarification	3658,0	95,4	-176,00	-
	Deep scarification	3446,0	89,8	-388,00	-
	Plowing	3593,0	93,7	-241,00	-

LSD 5% - 367,35; LSD 1% - 637,98; LSD 0,1% - 1298,81.

Analyzing the table of comparisons for the production obtained per unit area, it is also noted in the classical fertilization system and in the supplementary fertilization system the variant of soil preparation with the disk. Thus, in the classical fertilization system (NPK) the difference in production compared to the control (soil preparation with the hard disc) for the other three types of soil preparation is negative and varies between - 74.5 kg and - 241 kg/ha, and in the supplementary fertilization system the difference compared to the control reaches a negative maximum of -388 kg/ha. In both cases, the lowest production was obtained when preparation the soil by deep scarification and the best results when preparation the soil with the disc.

Regarding the main productivity elements: panicle length, number of grains/panicle, grain weight/panicle, straw length, thousand kernel weight (TKW), hectoliter weight (HLW) – these were obtained by analyzing a number of 50 plants for each experimental variant and are presented in the following figures.

The panicle length varies between 17.72 cm – 20.98 cm in the case of basic fertilization, respectively 18.42 cm – 22.78 cm in the case of nitrogen-supplemented fertilization.

The number of grains/panicle is higher in the variants worked by deep scarification and plowing, compared to those worked with the disk or by superficial scarification. The weight of grains/panicle, however, is higher in the case of the first two types of soil cultivation.




Figure 3. Evolution of productivity elements – straw lenght, panicle lenght, number of branches/ panicle




Figure 4. Evolution of productivity elements – number of grains/ panicle, grain weight/ panicle

And the hectoliter mass varies depending on the type of fertilization and the type of soil cultivation (Figure 5).

The thousand kernel weight when applying basic fertilization varies between 28.82 g – 29.98 g, with the highest value recorded in the plowing variant, closely followed by the other variants. When applying supplemented fertilization, TKW varies from 28.81 g to 29.21 g, with the highest value in the discing variant.

The hectoliter weight also varies depending on the type of fertilization and the type of soil cultivation (Figure 5).

In this case, the work of cultivating the soil with the disk is also noticeable, followed by the work of superficial scarification.




Figure 5. Evolution of productivity elements – TKW, HLW

Regarding the quality of the crop, the protein percentage was determined. The analysis was carried out in the quality laboratory of SCDA Lovrin, using the Kjeldahl method. The results obtained are presented in Table 2. The analysis of the presented data indicates a variation in the protein content in the range of 12.05% - 12.8% when applying basic fertilization and in the range of 13.35% - 14.05% when applying additional nitrogen fertilization.

Table 2. Comparison table for protein content - influence of fertilization and type of tillage

Experimental factors		Protein (%)	Difference	%	Significant
NPK	Disc harrowing	12.80	100.0	0.00	Control
	Light scarification	12.40	96.90	-0.40	-
	Deep scarification	12.40	96.90	-0.40	-
	Plowing	12.05	94.10	-0.75	00
NPK + NH ₄ NO ₃	Disc harrowing	14.05	100.0	0.00	Control
	Light scarification	13.60	96.80	-0.45	0
	Deep scarification	13.35	95.00	-0.70	00
	Plowing	13.40	95.40	-0.65	00

LSD 5% - 0.44; LSD 1% - 0.62; LSD 0,1% - 0.88.

The influence of additional nitrogen fertilization, as shown by the variance analysis (Table 3), is distinctly positive for the 1% transgression probability.

Table 3. Comparison table for protein content - influence of fertilization

Fertilization	Protein (%)	Difference	%	Significant
NPK	12.41	100.0	0.00	Control
NPK + NH ₄ NO ₃	13.60	109.6	1.19	**

LSD 5% - 0.45; LSD 1% - 1.05; LSD 0,1% - 3.33.

The type of tillage also has a significant influence on the quality of the spring oat crop (Table 4). Thus, compared to the minimal tillage with a heavy disc (control), the other three types of tillage significantly, distinctly significantly and very significantly influence this quality index analyzed.

Table 4. Comparison table for protein content - influence of type of tillage

Fertilization	Difference	%	Significant	Difference
Disc harrowing	13.43	100.0	0.00	Control
Light scarification	13.00	96.80	-0.43	0
Deep scarification	12.88	95.90	-0.55	00
Plowing	12.73	94.80	-0.70	000

LSD 5% - 0.31; LSD 1% - 0.44; LSD 0,1% - 0.62.

It is noted as in the case of production and productivity elements - the work of preparing the soil with the heavy disc.

CONCLUSIONS

Of the four types of soil tillage practiced for the establishment of spring straw cereal crops, for the western area of Romania the most efficient is the method of preparing the seedbed with the heavy disc. By using this method, the costs of land preparation are reduced by 73%. The highest production was recorded in the variant of soil cultivation with a heavy disk, both with basic fertilization and with the application of additional nitrogen fertilization, the production increase compared to the classic soil preparation work with a plow being 241 kg/ha. The formation of productivity elements and the quality of the crop is positively influenced by this type of soil cultivation.

The economic efficiency of soil tillage with the disc also derives from the higher yields per hectare obtained compared to the other three methods, the quality of the productions and a better storage and preservation of water in the soil.

ACKNOWLEDGMENTS

The present data were generated following the implementation of the ADER 1.2.3 project.

REFERENCES

1. Battisti DS., Naylor RL. Historical warnings of future food insecurity with unprecedented seasonal heat. *Science*, 2009, 323, pp 240–244.
2. Bărdaş M., Rusu T., ŞimonA., CheţanF., Popa A., Vâtcă S. Effect of the tillage systems and foliar fertilizations on assimilation, production and quality of wheat in the Transylvanian Plain conditions. *AgroLife Scientific Journal*, 2022, 11(2).
3. Byun YJ., Koo MY., JooHJ., Ha-Lee Y-M., Lee D-H. Comparative analysis of gene expression under cold acclimation, deacclimation and reacclimation in *Arabidopsis*. *Physiologia Plantarum*, 2014, 152, pp 256–274.
4. Ceclan A., ŞimonA., Russu F., Bărdaş M., CheţanF., Popa A. The influence of nitrogen, phosphorus, and potassium fertilizers on spring barley production and harvest increase. *Life Science and Sustainable Development Lovrin*, 3(2), p.60-66, LSSD -ISSN 2734 –5068.
5. Cheţan F., Cheţan C. Improvement of the water management by applying the no tillage system at the winter wheat at ARDS Turda. Symposium Life sciences for sustainable development. UASMV Cluj 26-28 September, Bulletin USAMV Cluj, 2021, 78(1), p: 22-27.
6. Cheţan F., CheţanC., MureşanuF., Suciul., Crişan I. Research regarding the efficiency of soybean cultivation in different tillage systems and their influence on soil compaction, accumulation and water storage. *Romanian Journal for Plant Protection*, 2021, 14,p.1-8. ISSN 2248 –129X; ISSN-L 2248 –129X
7. Chiriţă S., Urdă C., Şimon A., Cheţan F., Rusu T. Influence of technological factors on quality and some yield elements of Andrada winter wheat variety, cultivated in Transylvania Plain conditions. *LSSD Journal*, 2024, 5(1), LSSD -ISSN 2734 –5068.
8. Copcea Duma A., Mihaş C., Popa D. Studies regarding minimum soil tillage. *Scientific Papers Series Management, Economic Engineering in Agriculture and Rural Development*, 2018,18(1), p.153-156, Print ISSN 2284-7995, E-ISSN 2285-3952.
9. Derpsch, R.-Frontiers in Conservation Tillage and Advances in Conservation Practice. In: D.E.Stott, R.H.Mohtar and G.C.Steinhardt (eds), 2001, Sustaining the Global Farm. Selected papers from the 10th International Soil Conservation Organization Meeting held May 24 -29, 1999at Purdue University and the USDA-ARS National Soil Erosion Research Laboratory, p 248-254.
10. Deryng D., Conway D., Ramankutty N., Price J., Warren R. Global crop yield response to extreme heat stress under multiple climate change futures. *Environmental Research Letters*, 2014,9.
11. Diez JM., D'Antonio CM., Dukes JS. et al. Will extreme climatic events facilitate biological invasions? *Frontiers in Ecology and the Environment*, 2012, 10, pp 249–257.
12. Dinuţa I.C., Podea M.M., Marin D. I. The influence of the conservative tillage system on the pea crop in the pedoclimatic conditions of A.R.D.S. Piteşti. *Scientific Papers. Series A. Agronomy*, 2024, Vol. LXVII(1), p.57-65. ISSN 2285-5785; ISSN CD-ROM 2285-5793; ISSN Online 2285-5807; ISSN-L 2285-5785.
13. Gürsoy S. Soil Compaction Due to Increased Machinery Intensity in Agricultural Production: Its Main Causes, Effects and Management. In volume *Technology in Agriculture*, 2021, chapter 5, Doi: 10.5772/intechopen.98564.
14. Hoffman, L.A. World Production and Use of Oats, in *The Oat Crop*, edited by R.W. Welch, , London, 1995, pp. 34–61;

15. Hua J. From freezing to scorching, transcriptional responses to temperature variations in plants. *Current Opinion in Plant Biology*, 2009, 12, pp 568–573. doi:10.1016/j.pbi.2009.07.012.
16. Ipcc, Climate Change: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, 2014, Eds Core Writing Team, RK Pachauri, LA Meyer, IPCC: Geneva, Switzerland.
17. Jia Y., Ding Y., Shi Y., Zhang X., Gong Z., Yang S. The cbfs triple mutants reveal the essential functions of CBFs in cold acclimation and allow the definition of CBF regulons in Arabidopsis. *New Phytologist*, 2016, 212, pp 345–353.
18. Mickelbart MV., Hasegawa PM., Bailey-Serres J. Genetic mechanisms of abiotic stress tolerance that translate to crop yield stability. *Nature Reviews Genetics*, 2015, 16, pp 237–251. doi:10.1038/nrg3901.
19. Mileusnić Z., Saljnikov E., Radojevic R., Petrovic D.-Soil compaction due to agricultural machinery impact. *Journal of Terramechanics*. 2022, 100, 51-60, 10.1016/j.jterra.2021.12.002.
20. Niu S., Luo Y., Li D., Cao S., Xia J., Li J., Smith MD. Plant growth and mortality under climatic extremes: an overview. *Environmental and Experimental Botany*, 2014, 98, pp 13–19.
21. Rampino P., Mita G., Pataleo S., De Pascali M., Di Fonzo N., Perrotta C. Acquisition of thermotolerance and HSP gene expression in durum wheat (*Triticum durum* Desf.) cultivars. *Environmental and Experimental Botany*, 2009, 66, pp 257–264.
22. Rusu T., Moraru P.I., Ranta O., Drocas I., Bogdan I., Pop A.I., Sopterean M.L. No-Tillage and minimum tillage –their impact on soil compaction, water dynamics, soil temperature and production on wheat, maize and soybean crop. *Bulletin UASVM Agriculture*, 2011, 68(1), p.318-323. Print ISSN 1843-5246; Electronic ISSN 1843-5386.
23. Senthil-Kumar M., Srikanthbabu V., Mohan Raju B., Ganeshkumar N., Shivaprakash N., Udayakumar M. Screening of inbred lines to develop a thermotolerant sunflower hybrid using the temperature induction response (TIR) technique: a novel approach by exploiting residual variability. *Journal of Experimental Botany*, 2003, 54, pp 2569–2578.
24. Şimon A., Ceclan A., Russu F., Bârdaş M., Cheţan F., Popa A. The influence of N and P fertilization on winter wheat yield. *Life Science and Sustainable Development Lovrin*, 2022, 3(2), p.54-59, LSSD -ISSN 2734–5068.
25. Thomashow MF. Molecular basis of plant cold acclimation: insights gained from studying the CBF cold response pathway. *Plant Physiology*, 2010, 154, pp 571–577.
26. Witcombe J., Hollington P., Howarth C., Reader S., Steele KA. Breeding for abiotic stresses for sustainable agriculture. *Philosophical Transactions of the Royal Society of London*, 2008, Series B, 363, pp 703–716.
27. Zhang Y., Zhang L., Yang N., Huth N., Wang E., van der Werf W., Evers JB., Wang Q., Zhang D., Wang R., Gao H., Anten NPR. Optimized sowing time windows mitigate climate risks for oats production under cool semi-arid growing conditions. *Agricultural and Forest Meteorology*, 2019, 266–267, pp 184–197.